

# Humbert polynomials and functions in terms of Hermite polynomials

C. Cesarano, Dario Assante

Faculty of Engineering, International Telematic University UNINETTUNO, Rome, Italy

**Abstract—** By starting from the standard definitions of the incomplete two-variable Hermite polynomials, we propose non-trivial generalizations and we show some applications to the Bessel-type functions as the Humbert functions. We also present a generalization of the Laguerre polynomials in the same context of the incomplete-type and we use these to obtain relevant operational techniques for the Humbert-type functions.

**Keywords—** Hermite Polynomials, Laguerre polynomials, Generating Functions, Orthogonal Polynomials, Humbert polynomials, Humbert Functions, Bessel Functions.

## I. INTRODUCTION

It is possible to introduce a generalization of the Hermite polynomials which are a vectorial extension of the ordinary Kampé de Fériét one-variable Hermite polynomials [1]. We have indicated this class of the Hermite polynomials, of two-index and two-variable, by the symbol  $He_{m,n}(x, y)$ , and we stated their definition through the following generating function:

$$e^{-\hat{h}^t \hat{M} z - \frac{1}{2} \hat{h}^t \hat{M} h} = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} \frac{t^m u^n}{m! n!} He_{m,n}(x, y), \quad (1)$$

where:

$$\underline{z} = \begin{pmatrix} x \\ y \end{pmatrix} \text{ and } \underline{h} = \begin{pmatrix} t \\ u \end{pmatrix}$$

are two vectors of the space  $\mathbb{R}^2$  such that:  $t \neq u$ ,  $(|t|, |u|) < +\infty$ , and the superscript “ $r$ ” denotes transpose.

A different generalization of the Hermite polynomials could be obtained by using the slight similar procedure onto the two-variable generalized Hermite polynomials [1,2]:

$$H_n(x, y) = n! \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \frac{y^r x^{n-2r}}{r!(n-2r)!}, \quad (2)$$

defined by the generating function of the form:

$$\exp(xt + yt^2) = \sum_{n=0}^{+\infty} \frac{t^n}{n!} H_n(x, y). \quad (3)$$

Let  $u$  and  $v$  continuous variables, such that  $u \neq v$  and  $(|u|, |v|) < +\infty$ ,  $\tau \in \mathbb{R}$ , we will say incomplete 2-dimensional Hermite polynomials, the polynomials defined by following generating function:

$$\exp(xu + yv + \tau uv) = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} \frac{u^m v^n}{m! n!} h_{m,n}(x, y | \tau). \quad (4)$$

This class of Hermite polynomials has been deeply studied for its importance in applications, as quantum mechanical problems, harmonic oscillator functions and also to investigate the statistical properties of chaotic light [3]. By using the techniques of the generating function method [4,5], it is easy to obtain the explicit form of the above polynomials:

$$h_{m,n}(x, y | \tau) = m! n! \sum_{r=0}^{[m,n]} \frac{\tau^r x^{m-r} y^{n-r}}{r!(m-r)!(n-r)!}, \quad (5)$$

where  $[m, n] = \min(m, n)$ .

An interesting particular case of this class of Hermite polynomials is presented when  $x = y = 1$  and  $\tau = x$ :

$$h_{m,n}(1, 1 | x) = g_{m,n}(x). \quad (6)$$

It is significant to study the polynomials  $g_{m,n}(x)$  since they can be used to define other forms of the incomplete 2-dimensional Hermite polynomials of the type  $h_{m,n}(\cdot, \cdot | \cdot)$  themselves and since they often appear in the description of the applications in quantum optics. From the relation (6) and by using the definitions (4) and (5), we can immediately write the following general relation:

$$h_{m,n}(x, y | \tau) = x^m y^n g_{m,n} \left( \frac{\tau}{xy} \right). \quad (7)$$

The incomplete 2-dimensional Hermite polynomials can be used to obtain different forms of the multi-index Bessel functions, in particular for the case of the Humbert functions. We remind that the ordinary cylindrical Bessel functions [6] are specified by the generating function:

$$\exp\left[\frac{x}{2}\left(t - \frac{1}{t}\right)\right] = \sum_{n=-\infty}^{+\infty} t^n J_n(x), \quad (8)$$

and a generalization of them, it is represented by the case of two-index, one-variable type [7,8]:

$$J_{m,n}(x) = \sum_{s=-\infty}^{+\infty} J_{m-s}(x) J_{n-s}(x) J_s(x), \quad (9)$$

with the following generating function:

$$\begin{aligned} \exp\left[\frac{x}{2}\left(u - \frac{1}{u}\right) + \left(v - \frac{1}{v}\right)\left(uv - \frac{1}{uv}\right)\right] = \\ = \sum_{m=-\infty}^{+\infty} \sum_{n=-\infty}^{+\infty} u^m v^n J_{m,n}(x), \end{aligned} \quad (10)$$

where  $x \in \mathbb{R}$  and  $u, v \in \mathbb{R}$ , such that  $0 < |u| \neq |v| < +\infty$ .

This class of Bessel functions satisfied analogous interesting relations as the ordinary Bessel functions. For instance, by deriving in the equation (10) with respect to  $x$ , we have:

$$\begin{aligned} \frac{1}{2}\left(u - \frac{1}{u}\right) + \left(v - \frac{1}{v}\right)\left(uv - \frac{1}{uv}\right) \sum_{m=-\infty}^{+\infty} \sum_{n=-\infty}^{+\infty} u^m v^n J_{m,n}(x) = \\ = \sum_{m=-\infty}^{+\infty} \sum_{n=-\infty}^{+\infty} u^m v^n \frac{d}{dx} J_{m,n}(x), \end{aligned} \quad (11)$$

which allows us to state the following recurrence relation:

$$\begin{aligned} \frac{d}{dx} J_{m,n}(x) = \frac{1}{2} \{ [J_{m-1,n}(x) - J_{m+1,n}(x)] + [J_{m,n-1}(x) - J_{m,n+1}(x)] + \\ + [J_{m-1,n-1}(x) - J_{m+1,n+1}(x)] \}. \end{aligned} \quad (12)$$

By using the same procedure, it is easy to obtain the other two recurrence relations for this class of Bessel functions:

$$\begin{aligned} \frac{2m}{x} J_{m,n}(x) = [J_{m,n-1}(x) - J_{m,n+1}(x)] + \\ + [J_{m-1,n-1}(x) - J_{m+1,n+1}(x)], \end{aligned} \quad (13)$$

and:

$$\begin{aligned} \frac{2n}{x} J_{m,n}(x) = [J_{m,n-1}(x) - J_{m,n+1}(x)] + \\ + [J_{m-1,n-1}(x) - J_{m+1,n+1}(x)]. \end{aligned} \quad (14)$$

It is interesting to note that, for  $x = 0$ , from the explicit form of the generalized two-index Bessel function (eq. (9)), we get:

$$J_{m,n}(0) = \sum_{s=-\infty}^{+\infty} J_{m-s}(0) J_{n-s}(0) J_s(0), \quad (15)$$

and, since:

$$J_s(0) \neq 0, \text{ when } s = 0, \quad (16)$$

we, finally, obtain:

$$J_{m,n}(0) = \delta_{m,0} \delta_{n,0}. \quad (17)$$

As a particular case of the two-index, one-variable Bessel functions, we can introduce the Humbert functions [9], by setting:

$$b_{m,n}(x, y | \tau) = \sum_{r=0}^{+\infty} \frac{\tau^r x^{m+r} y^{n+r}}{r!(m+r)!(n+r)!}, \quad (18)$$

defined through the following generating function:

$$\exp(xu + yv + \tau uv) = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} u^m v^n b_{m,n}(x, y | \tau). \quad (19)$$

It is evident the similar structure between these functions and the incomplete 2-dimensional Hermite polynomials presented previously (see eqs. (4,5)). For this reason, the Humbert functions are usual exploited in connection with the Hermite polynomials of the type  $h_{m,n}(x, y | \tau)$ .

We can immediately note, for instance, that the Humbert functions could be expressed in terms of the incomplete Hermite polynomials. By rewriting, in fact, the expression in equation (19), we have:

$$\exp\left[xu + yv + \tau uv - \tau\left(uv - \frac{1}{uv}\right)\right] = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} u^m v^n b_{m,n}(x, y | \tau), \quad (20)$$

and, from the generating function of the ordinary Bessel function (eq. (8)), we find:

$$b_{m,n}(x, y | \tau) = \sum_{r=0}^{[m,n]} \frac{(-1)^r h_{m-r,n-r}(x, y | \tau) J_r(2\tau)}{r!(m-r)!(n-r)!}. \quad (21)$$

In the following, we will indicate with:

$$g_{m,n}(x) \text{ and } b_{m,n}(x),$$

the Humbert polynomials and the Humbert functions respectively. In the next sections we will study the properties of these particular polynomials and functions and we will see some their non trivial generalizations along with the analysis of the related applications to facilitate some operational computation.

## II. RELEVANT PROPERTIES FOR HUMBERT POLYNOMIALS AND FUNCTIONS

In the previous section we have introduced the incomplete 2-dimensional Hermite polynomials through the relations (4,5). By using the equivalences stated in equations (6,7), we can now state the expression of the generating function for the Humbert polynomials  $g_{m,n}(x)$ . We have:

$$\exp(u + v + xuv) = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} \frac{u^m v^n}{m! n!} g_{m,n}(x) \quad (22)$$

where, again,  $u \neq v$  and  $(|u|, |v|) < +\infty, \tau \in \mathbb{R}$ .

By following the same procedure used to derive the recurrence relations related to the two-index, one-variable Bessel function in the previous section, we can find similar expressions for this class of Humbert polynomials. In fact, by deriving, respectively, with respect to  $x, u$  and  $v$ , we obtain:

$$\begin{aligned} \frac{d}{dx} g_{m,n}(x) &= mng_{m-1,n-1}(x), \\ g_{m+1,n}(x) &= g_{m,n}(x) + mxg_{m-1,n}(x), \\ g_{m,n+1}(x) &= g_{m,n}(x) + nxg_{m,n-1}(x). \end{aligned} \quad (23)$$

By using the above relations, it is possible to state the differential equation satisfied by the polynomials  $g_{m,n}(x)$ . After an easy manipulation of the equations in the (23), we get:

$$mg_{m-1,n}(x) = \left(m - x \frac{d}{dx}\right) g_{m,n}(x), \quad (24)$$

$$ng_{m,n-1}(x) = \left(n - x \frac{d}{dx}\right) g_{m,n}(x) \quad (25)$$

and also:

$$mng_{m-1,n-1}(x) = \left(m - x \frac{d}{dx}\right) \left(n - x \frac{d}{dx}\right) g_{m,n}(x). \quad (26)$$

After equating equation (24) with the first of the relations obtained in (23), we can state the following differential equation solved by the Humbert polynomials:

$$x^2 y'' - [(m+n-1)x+1]y' + mny = 0. \quad (27)$$

We note that, from the equation (25), it also follows:

$$\begin{aligned} g_{m+1,n}(x) &= g_{m,n}(x) + nxg_{m,n}(x) - x^2 \frac{d}{dx} g_{m,n}(x), \\ g_{m,n+1}(x) &= g_{m,n}(x) + mxg_{m,n}(x) - x^2 \frac{d}{dx} g_{m,n}(x) \end{aligned} \quad (28)$$

Which suggest the introduction of the following operators:

$$\begin{aligned} \hat{S}_m^+ &= 1 + \hat{n}x - x^2 \frac{d}{dx}, \\ \hat{S}_n^+ &= 1 + \hat{m}x - x^2 \frac{d}{dx}, \end{aligned} \quad (29)$$

where we have denoted with the symbols:

$$\hat{m} \text{ and } \hat{n}$$

a kind of number operators, in the sense that their action read as following:

$$\hat{m} \hat{n} g_{s,r}(x) = sr g_{s,r}(x).$$

It is now evident, by using the relations stated in the equations (23-28) and by the definition of the operators expressed in equation (29), that the following expressions hold:

$$\begin{aligned} \hat{S}_m^+ g_{m,n}(x) &= g_{m+1,n}(x), \\ \hat{S}_n^+ g_{m,n}(x) &= g_{m,n+1}(x). \end{aligned} \quad (30)$$

The above relations, combined with the first in the equation (23), allow us to state the following relevant differential equation:

$$\begin{aligned} \frac{d}{dx} \left[ 1 + (n+1)x - x^2 \frac{d}{dx} \right] \left[ 1 + mx - x^2 \frac{d}{dx} \right] g_{m,n}(x) &= \\ = (m+1)(n+1)g_{m,n}(x). \end{aligned} \quad (31)$$

It is possible to derive similar relations regarding the Humbert functions. Before to proceed, we remind that, the function defined by the following generating function:

$$\exp\left(t - \frac{x}{t}\right) = \sum_{n=-\infty}^{+\infty} t^n C_n(x) \quad (32)$$

is known as the Tricomi function [10], which its explicit form is:

$$C_n(x) = \sum_{r=0}^{+\infty} \frac{(-1)^r x^r}{r!(n+r)!}. \quad (33)$$

It is possible to introduce a generalization of the above function in the sense of the Humbert functions. In fact, from the equation (18) it is immediately recognized that we can call generalized Tricomi function, the function expressed by the following relation:

$$C_{m,n}(x) = b_{m,n}(1,1|x). \tag{34}$$

By using the same procedure outlined above, we can derive, for the Humbert functions the analogous recurrence relations stated for the polynomials  $g_{m,n}(x)$ . In fact, by considering the relation (34), we have:

$$\begin{aligned} \frac{d}{dx} C_{m,n}(x) &= C_{m+1,n+1}(x), \\ mC_{m,n}(x) &= C_{m-1,n}(x) - xC_{m+1,n+1}(x), \\ nC_{m,n}(x) &= C_{m,n-1}(x) - xC_{m+1,n+1}(x). \end{aligned} \tag{35}$$

We can combine the above relations, to get:

$$\begin{aligned} C_{m-1,n}(x) &= \left(m + x \frac{d}{dx}\right) C_{m,n}(x), \\ C_{m,n-1}(x) &= \left(n + x \frac{d}{dx}\right) C_{m,n}(x). \end{aligned} \tag{36}$$

These last relations suggest to introduce similar operators acting on these generalized Tricomi function as well as we have done for the Humbert polynomials. We have indeed:

$$\begin{aligned} \hat{E}_m^- &= m + x \frac{d}{dx}, \\ \hat{E}_n^- &= n + x \frac{d}{dx}, \\ \hat{E}_{m,n}^{+,+} &= \frac{d}{dx}. \end{aligned} \tag{37}$$

We have used, again, the same notation as expressed for the operators in equation (29). By following the same procedure used for the Humbert polynomials, we can easily to state the following differential equation:

$$x^2 y''' - (m+n+3)xy'' + (mn+m+n+1)y' = y. \tag{38}$$

### III. FURTHER GENERALIZATIONS FOR HUMBERT POLYNOMIALS AND FUNCTIONS AND INCOMPLETE LAGUERRE POLYNOMIALS

In the paper [10], we have showed some relations linked the cylindrical Bessel function and the Tricomi function; in particular, we have seen that:

$$x^{-\frac{n}{2}} J_0(2\sqrt{x}) = C_n(x) = \sum_{r=0}^{+\infty} \frac{(-1)^r x^r}{r!(n+r)!}. \tag{39}$$

The relations stated in the previous sections and the structure of the  $O^h$  order Tricomi function, allow us to introduce a generalization of the Laguerre polynomials.

We will say incomplete 2-dimensional Laguerre polynomials, the polynomials defined by the following generating function:

$$\exp(u+v)C_0(xuv) = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} \frac{u^m v^n}{m! n!} l_{m,n}(x), \tag{40}$$

where their explicit form reads:

$$l_{m,n}(x) = m!n! \sum_{r=0}^{[m,n]} \frac{(-1)^r x^r}{(r!)^2 (m-r)!(n-r)!}. \tag{41}$$

It is evident the similar structure with the Humbert polynomials discussed in the previous sections.

We remind that the ordinary Laguerre polynomials [10] have the following operational expression:

$$L_n(x) = \left(1 - \hat{D}^{-1}\right)^n, \tag{42}$$

where  $\hat{D}_x^{-1}$  denotes the inverse of the derivative operator [11-14], being essentially an integral operator, it will be specified by the operational rule:

$$\hat{D}_x^{-1}(1) = \frac{x^n}{n!}. \tag{43}$$

From the above considerations, we can firstly write the following expression for the  $O^h$  Tricomi function:

$$C_0(x) = \sum_{r=0}^{+\infty} \frac{(-1)^r \hat{D}_x^{-r}}{r!} = \exp\left(-\hat{D}_x^{-1}\right), \tag{44}$$

and then we can state the important equivalence between the Humbert polynomials and the incomplete 2-dimensional Laguerre polynomials, that is:

$$l_{m,n}(x) = g_{m,n}\left(-\hat{D}_x^{-1}\right). \tag{45}$$

By recalling the explicit forms of the generalized two-variable Laguerre polynomials:

$$L_n(x, y) = \left( y - D_x^{-1} \right)^n = n! \sum_{r=0}^n \frac{(-1)^r y^{n-r} x^r}{(r!)^2 (n-r)!} \quad (46)$$

and from the particular expression of their generating function, in terms of the Tricomi function:

$$\sum_{n=0}^{+\infty} \frac{t^n}{n!} L_n(x, y) = \exp(yt) C_0(xt), \quad (47)$$

we can finally establish a link between the incomplete 2-dimensional laguerre polynomials and the generalized Laguerre of the form  $L_n(x, y)$ . We have:

$$l_{m,n}(x) = m! n! \sum_{r=0}^{[m,n]} \frac{g_{m-r,n-r}(-y) L_r(x, y)}{r!(m-r)!(n-r)!}. \quad (48)$$

The considerations and the following results obtained to define the incomplete 2-dimensional polynomials, can be used to introduce a similar generalization for the Humbert functions. By considering indeed the following generating function:

$$\exp(u+v) C_0\left(\frac{x}{uv}\right) = \sum_{m=0}^{+\infty} \sum_{n=0}^{+\infty} u^m v^n A_{m,n}(x), \quad (49)$$

we easily obtain the explicit form of the function  $A_{m,n}(x)$ :

$$A_{m,n}(x) = \sum_{r=0}^{+\infty} \frac{(-1)^r x^r}{(r!)^2 (m+r)!(n+r)!}. \quad (50)$$

It is easy to note the analogy between the above expression and the generalized Tricomi function presented in the previous section. We find in fact:

$$A_{m,n}(x) = C_{m,n}\left(-D_x^{-1}\right). \quad (51)$$

In the same way, it is possible obtain an expression of the functions  $A_{m,n}(x)$  involving the generalized two-variable Laguerre polynomials. From the relation stated in equation (48) and from the (50), we have:

$$A_{m,n}(x) = \sum_{r=0}^{+\infty} \frac{g_{m+r,n+r}(-y) L_r(x, y)}{r!(m+r)!(n+r)!}. \quad (52)$$

#### IV. CONCLUDING REMARKS

Before closing the paper, we want just to mention how the concepts and the formalism discussed in the previous sections allows also the generalizations of other simple distribution functions like the Poisson distribution.

By using the Tricomi function of order  $m$ :

$$C_m(-x) = \sum_{r=0}^{+\infty} \frac{x^r}{r!(m+r)!}, \quad (53)$$

we can indeed define the following two-index distribution:

$$P_n(x; m) = \frac{x^n}{n!(m+n)! C_m(-x)}, \quad (54)$$

where the generating function is given by the relation:

$$\frac{C_m(-xt)}{C_m(-x)} = \sum_{n=0}^{+\infty} \frac{t^n}{n!} P_n(x; m). \quad (55)$$

The evaluation of the associated momenta can be easily simplified with the use of the well known property, satisfied by the Tricomi functions:

$$(-1)^r \left( \frac{d}{dx} \right)^r C_n(x) = C_{n+r}(x). \quad (58)$$

Accordingly, we calculate the following average values:

$$\langle n \rangle = \frac{C_{m+1}(-x)}{C_m(-x)}, \quad (59)$$

$$\langle n^2 \rangle = \frac{C_{m+2}(-x)}{C_m(-x)} + \langle n \rangle.$$

The higher order moments are also given by similar closed relations.

It is remarkable about this probability distribution that, unlike the Poisson distribution, the variance:

$$\sigma = \sqrt{n^2 - n}, \text{ where } \bar{a} = \langle a \rangle \quad (60)$$

is smaller than  $\sqrt{\bar{a}}$ .

This type of distribution can be exploited in quantum mechanics within the context of bunching phenomena. This example show that the use of multi-index polynomials and Bessel-type functions with their associated formalism offers wide possibilities in the applications of pure and applied mathematics.

#### REFERENCES

- [1] Appell, P., Kampé de Fériet, J., "Fonctions hypergéométriques et hypersphériques. Polynomes d'Hermite", *Gauthier-Villars*, Paris, 1926.
- [2] Dattoli, G., "Incomplete 2-D Hermite polynomials: Properties and applications", *Journal of Mathematical Analysis and Applications*, 284 (2), pp 447-454, 2003.
- [3] Dodonov, V.V., Man'ko, V.I., "New relations for two-dimensional Hermite polynomials", *Journal of mathematical Physics*, 35 (8), pp. 4277-4294, 1994.

- [4] Gould, H.W., Hopper, A.T., "Operational formulas connected with two generalizations of Hermite polynomials", *Duke Math. J.*, 29, pp. 51-62, 1962.
- [5] Srivastava, H.M., Manocha, H.L., "A treatise on generating functions", Wiley, New York, 1984.
- [6] Watson, J.H., "A treatise on Bessel functions", *Cambridge University press*, 1958.
- [7] Dattoli, G., Torre, A. "Theory and applications of generalized Bessel functions", *Aracne, Rome*, 1996.
- [8] Cesarano, C., Assante, D., "A note on generalized Bessel functions", *Int. J. of Math. Models and Methods in Appl. Sci.*, 7 (6), pp 625-629, 2013.
- [9] Aktas, R., Sahin, R., Altin, A., "On a multivariable extension of the Humbert polynomials", *Applied Mathematics and Computation*, 218 (3), pp. 662-666, 2011.
- [10] Cesarano, C., "Monomiality Principle and related operational techniques for Orthogonal Polynomials and Special Functions", *Int. J. of Math. Models and Methods in Appl. Sci.*, to appear, 2013.
- [11] Dattoli, G., Lorenzutta, S., Ricci, P.E., Cesarano, C., "On a family of hybrid polynomials", *Integral Transforms and Special Functions*, 15 (6), pp. 485-490, (2004).
- [12] Dattoli, G., Lorenzutta, Cesarano, C., "Bernstein polynomials and operational methods", *Journal of Computational Analysis and Applications*, 8 (4), pp. 369-377, 2006.
- [13] Dattoli, G., Srivastava, H.M., Cesarano, C., "The Laguerre and Legendre polynomials from an operational point of view", *Applied Mathematics and Computation*, 124 (1), pp. 117-127 (2001).
- [14] Dattoli, G., Cesarano, C., "On a new family of Hermite polynomials associated to parabolic cylinder functions", *applied Mathematics and Computation*, 141 (1), pp. 143-149, 2003.

**Clemente Cesarano** is assistant professor of Mathematical Analysis at Faculty of Engineering- International Telematic University UNINETTUNO-Rome, ITALY. He is coordinator of didactic planning of the Faculty and he also is coordinator of research activities of the University. Clemente Cesarano is Honorary Research Associates of the Australian Institute of High Energetic Materials His research activity focuses on the area of Special Functions, Numerical Analysis and Differential Equations.

He has work in many international institutions as ENEA (Italy), Ulm University (Germany), Complutense University (Spain) and University of Rome La Sapienza (Italy). He has been visiting researcher at Research Institute for Symbolic Computation (RISC), Johannes Kepler University of Linz (Austrian). He is Editorial Board member of the Research Bulletin of the Australian Institute of High Energetic Materials, the Global Journal of Pure and Applied Mathematics (GJPAM), the Global Journal of Applied Mathematics and Mathematical Sciences (GJ-AMMS), the Pacific-Asian Journal of Mathematics, the International Journal of Mathematical Sciences (IJMS), the Advances in Theoretical and Applied Mathematics (ATAM), the International Journal of Mathematics and Computing Applications (IJMCA). Clemente Cesarano has published two books and over than fifty papers on international journals in the field of Special Functions, Orthogonal Polynomials and Differential Equations.

## Creative Commons Attribution License 4.0 (Attribution 4.0 International, CC BY 4.0)

This article is published under the terms of the Creative Commons Attribution License 4.0

[https://creativecommons.org/licenses/by/4.0/deed.en\\_US](https://creativecommons.org/licenses/by/4.0/deed.en_US)